

Medication Tracker

NUCALA works over time, which is why it's important to track your ongoing treatments with your doctor. Working your upcoming appointments into your routine and managing your medication changes can require some extra organization.

NUCALA is an add-on, prescription maintenance treatment for patients 12 and older with severe eosinophilic asthma. NUCALA is not used to treat sudden breathing problems.

This tool will provide a place to write down your current medications and dosages, as well as keep track of dosage adjustments. As you continue taking NUCALA, this tool can help you and your doctor see any changes in your medications over time.

	1st Injection		Ongoing Treatment							
	Start Date ____/____/____		Month 3 ____/____/____		Month 6 ____/____/____		Month 9 ____/____/____		Month 12 ____/____/____	
Current Medications	Dose	Frequency	Dose	Frequency	Dose	Frequency	Dose	Frequency	Dose	Frequency

Important Safety Information

Do not use NUCALA if you are allergic to mepolizumab or any of the ingredients in NUCALA. Do not use to treat sudden breathing problems.

Please see additional Important Safety Information on Page 2.





Remember: Your doctor may reduce your use of oral steroids, like prednisone, after you start NUCALA. Remember to always consult your doctor before adjusting your medications. Your results may vary.

Important Safety Information (cont'd)

NUCALA can cause serious side effects, including:

- **allergic (hypersensitivity) reactions, including anaphylaxis.** Serious allergic reactions can happen after you get your injection of NUCALA. Allergic reactions can sometimes happen hours or days after you get a dose of NUCALA. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction:
 - swelling of your face, mouth, and tongue
 - breathing problems
 - fainting, dizziness, feeling light-headed (low blood pressure)
 - rash
 - hives
- **Herpes zoster infections** that can cause shingles have happened in people who received NUCALA.

Before receiving NUCALA, tell your healthcare provider about all of your medical conditions, including if you:

- are taking oral or inhaled corticosteroid medicines. **Do not** stop taking your other asthma medicines, including your corticosteroid medicines, unless instructed by your healthcare provider because this may cause other symptoms to come back.
- have a parasitic (helminth) infection.
- are pregnant or plan to become pregnant. It is not known if NUCALA may harm your unborn baby.
 - A pregnancy registry for women who receive NUCALA while pregnant collects information about the health of you and your baby. You can talk to your healthcare provider about how to take part in this registry or you can get more information and register by calling 1-877-311-8972 or visit www.mothersbaby.org/asthma.
- are breastfeeding or plan to breastfeed. You and your healthcare provider should decide if you will use NUCALA and breastfeed. You should not do both without talking with your healthcare provider first.
- are taking prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of NUCALA include: headache, injection site reactions (pain, redness, swelling, itching, or a burning feeling at the injection site), back pain, and weakness (fatigue).

Please see additional Important Safety Information on Page 1.

Please see full Prescribing Information, including Patient Information Leaflet, for NUCALA available at NUCALA.com.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Trademarks are owned by or licensed to the GSK group of companies.



©2018 GSK group of companies or its licensor.
Printed in USA. 823267R0 January 2018

Nucala 
(mepolizumab)
for Subcutaneous Injection
100 mg/vial